arXiv:1604.00341v1 [math.GR] 1 Apr 2016

April 4, 2016

CONSTRUCTION OF RIGHT GYROGROUP STRUCTURES ON A NON-ABELIAN GROUP

AKHILESH CHANDRA YADAV

Abstract. In this article we gave the notion of particular type of functions on a group termed as class assigned functions . Using class assigned functions, we have constructed several right gyrogroup structures on a given non-abelian group.

Keywords: Right transversals, Right loops, Gyrotransversals.
MSC: 20N05.

1. Introduction

A gyrogroup is a grouplike structure that is defined in [3] along with a weaker structure called a left gyrogroup. In [4], Foguel and Ungar proved that any given group can be turned into a left gyrogroup which is, in turn, a gyrogroup if and only if the given group is central by a $2-Engel$ group. The importance of left gyrogroups stems from the facts that (i) any gyrotransversal groupoid is a left gyrogroup, and (ii) any left gyrogroup is a twisted subgroup in a specifed group. Let K be a group and let G be the inner product of the group K by $Inn(K)$, where $Inn(K)$ is the inner automorphism group of K whose generic element α_k denotes conjugation by $k \in K$. Then, the diagonal transversal D generated by K (in G) is a subset of G given by

$$D = \{(k; \alpha_k) \mid k \in K\}.$$

Then the diagonal transversal D is a gyrotransversal (Theorem 3.2, [4]). Indeed this operation induces a binary operation $\odot$ on K so that the groupoid $(K, \odot)$ is a left gyrogroup (see, [4]). The operation $\odot$ on K is given by $x \odot y = x^2yx^{-1}$. It is convenient for us to study right transversals and right conjugation action. Thus, in our terminology, $(K, \odot)$ is a right gyrogroup, where $x \odot y = y^{-1}xy^2$. In this article, we construct several right gyrogroup structures on a non-abelian group.

2. Preliminaries

Let S be a right transversal to the subgroup H in a group G. Let $g : S \to H$ be a map with $g(e) = 1$, where e denotes the identity of S with respect to transversal operation and 1 denotes the identity of

group H. Consider the right transversal $S_g = \{g(x)x \mid x \in S\}$ to H. Then

Proposition 2.1. [2] *The right quasigroup (S_g, o') is isomorphic to the right quasigroup (S, o_g) where o_g is the operation given by $xo_gy = x\theta g(y)oy$.*

The operation o_g is known as the deformed operation on S.

Definition 2.2. [2] *Let (S, o) be a magma together with a right identity e and together with a right inverse a' for every element $a \in S$ with respect to e in the sense that $aoa' = e$. Then (S, o) will be called a right gyrogroup if*

(1) *for any $x, y, z \in S$ there exists a unique element $x\theta f(y, z) \in S$ such that*

$$(xoy)oz = x\theta f(y, z)o(yoz), \tag{2.1}$$

(2) *the map $f(y, z) : S \to S$ given by $f(y, z)(x) = x\theta f(y, z)$ is an automorphism of (S, o) and*
(3) *for all $y \in S$,*

$$f(y, y') = I_S. \tag{2.2}$$

Definition 2.3. [2] *A right transversal S to a subgroup H with identity e in a group G is called a gyrotransversal if $S = S^{-1}$ and $h^{-1}xh \in S$, $\forall\ x \in S$ and $h \in H$.*

Proposition 2.4 ([2], Representation theorem for right gyrogroups)**.** *A right loop (S, o) is a right gyrogroup if and only if it is a gyrotransversal to G_S in its group extension G_SS.*

In [2], we have shown the following:

Lemma 2.5. [2] *Let S be a gyrotransversal to a subgroup H in a group G and g be a map from S to H such that $g(e) = e$. Then the transversal $S_g = \{g(x)x \mid x \in S\}$ is a gyrotransversal if and only if*

$$g(x^{-1}) = g(x)^{-1} \tag{2.3}$$

and g is equivariant in the sense that

$$g(h^{-1}xh) = h^{-1}g(x)h \tag{2.4}$$

for all $x \in S$ and $h \in H$.

3. Construction of Right Gyrogroups

Definition 3.1. *Let G be a given non-abelian group. Define a binary operation $\cdot$ on $G \times G$ as*

$$(a, x) \cdot (b, y) = (ab[b^{-1}xb, y^{-1}], b^{-1}xboy),$$

where $xoy = y^{-1}xy^2$. Then $(G \times G, \cdot)$ is a group. The identity of the group $G \times G$ is $(1, 1)$, where 1 denotes the identity of group G and the inverse of (a, x) is $(a^{-1}, a^{-1}x^{-1}a)$, for each (a, x) in $G \times G$.

One may easily observe that the group $G \times G$ contains $G \times \{1\} (\simeq G)$ as a subgroup and $\{1\} \times G$ as a gyrotransversal to the subgroup $G \times \{1\}$ in the group $G \times G([1,\, 2])$.

Proposition 3.2. *Let $n \in \mathbb{N}$ and g be a map from a group G into itself with $g(1) = 1$ and $g(x) = x^n$ for each $x \in G$. Then $S_g = \{(g(x), x) \,|\, x \in G\}$ is a gyrotransversal to the subgroup $H(= G \times \{1\})$ in $G \times G$.*

Proof: Since $(x^{-1})^n = (x^n)^{-1}$ and $(h^{-1}xh)^n = h^{-1}x^nh$ for all $x, h \in G$. Thus the result follows from Lemma 2.5. □

Let G be a group. Let R be a relation on G given by aRb if either $a = b$ or $a = b^{-1}$ or $a \sim b$ or $a \sim b^{-1}$, where $\sim$ is a conjugacy relation on group G. Then R is an equivalence relation on G. The equivalence class of $w \in G$ is denoted by $[w]_R$.

Definition 3.3. *Let G be a group. A function $k : G \to \mathbb{N} \cup \{0\}$ is called a class assigned function on G if $k([1]_R) = 0$ and k is constant on each class $[w]_R$.*

Proposition 3.4. *Let k be a class assigned function on a free group F. Then it determines a function $g : F \to F$ given by $g(w) = w^{k(w)}$ so that the corresponding transversal $S_g = \{(g(w), w) \,|\, w \in F\}$ to the subgroup $H = F \times \{1\}$ in $F \times F$ is a gyrotransversal.*

Proof: By definition of k, $k(w) = k(w^{-1}) = k(u^{-1}wu)$, for all u, w in F. Therefore $g(w^{-1}) = g(w)^{-1}$ and $g(u^{-1}wu) = (u^{-1}wu)^{k(u^{-1}wu)} = u^{-1}w^{k(u^{-1}wu)}u = u^{-1}w^{k(w)}u = u^{-1}g(w)u$. By Lemma 2.5, S_g is a gyrotransversal. □

Proposition 3.5. *Let $g : F \to F$ be a map satisfying the property that*
(i). $g(1) = 1$,
(ii). $g(w^{-1}) = g(w)^{-1}$,
(iii). $g(u^{-1}wu) = u^{-1}g(w)u$, for all $u, w \in F$.
Then g determines a class assigned function k on free group F such that $g(w) = w^{k(w)}$.

PROOF: For $w \in F \setminus \{1\}$, (iii) gives $g(w) = w^{-1}g(w)w$. Thus $g(w)$ belongs to the centralizer of w. But F is a free group, therefore $g(w) = w^{k(w)}$, for unique $k(w) \in \mathbb{N} \cup \{0\}$. Then (ii) and (iii) implies that $k(w^{-1}) = k(w)$ and $k(u^{-1}wu) = k(w)$ for all $u, w \in F$. Define $k(1) = 0$, thus $k(w)$ determines a class assigned function k. □

Corollary 3.6. *There is a bijective correspondence between the set of all class assigned functions and the set of all maps g described as in the Prop. (3.5). In particular, there is a bijective correspondence between the set of all class assigned functions and the set of all gyrotransversals to the subgroup H in $F \times F$.*

Theorem 3.7. *Let G be a non-abelian group and k be a class assigned function on G. Then it determines a function $g : G \to G$ given by $g(w) = w^{k(w)}$ such that S_g is a gyrotransversal to $G \times \{1\}$ in $G \times G$. Define a binary operation o_k on G by*

$$xo_k y = y^{-k(y)}xy^{k(y)+1}. \tag{3.1}$$

Then (G, o_k) is a right gyrogroup.

PROOF: By the property of class assigned function k, g satisfies $g(x^{-1}) = g(x)^{-1}$ and $g(h^{-1}xh) = h^{-1}g(x)h$ for all x, h in G. Thus S_g is a gyrotransversal to $G \times \{e\}$ in the group $G \times G$ and so deformed operation o_k on G is given by Equation 3.1. Observed that $y^{-1}o_k y = e = yo_k y^{-1}$. By a little computation the map $f(y, z)(x)$ determined by the equation $(xoy)oz = f(y, z)(x)o(yoz)$ is $f(y, z)(x) = x\theta\left[y^{k(y)}(yz)^{-k(yo_k z)}z^{k(z)}\right]$, where $x\theta a = a^{-1}xa$ for all $x, a \in G$. Clearly $f(y, z)$ is an inner automorphism of (G, o_k). Also $f(y, y^{-1})(x) = x = f(y^{-1}, y)(x)$ and so $f(y, y^{-1}) = I_G$. Thus, (G, o_k) is a right gyrogroup. □

Remark 3.8. *We do not know if its converse true.*

Example 3.2. *Take $G = S_3$, the symmetric group of degree 3. Let k be a class assigned function on S_3.*

If $k(x) = 0$ for all x in S_3. Then the corresponding right gyroroup (S_3, o_k) is a group S_3 itself.

If $k(I) = 0, k(x) = 1$ for all $x \in S_3 \setminus \{I\}$. Then the group generated by automorphisms $f(y, z)$ is isomorphic to $[S_3, S_3] = A_3$.

If $k(I) = 0$, $k((12))$=$k((2\ 3))$= $k((1\ 3))$=1, $k((1\ 2\ 3))$= $k((1\ 3\ 2))$= 2. Then $f((12), (132))(x) = x\theta(13)$, $f((12), (123))(x) = x\theta(23)$ and $f((23), (132))(x) = x\theta(12)$. Thus, (S_3, o_k) is a right gyrogroup for which the group generated by automorphisms $f(y, z)$ is isomorphic to S_3.

The above right gyrogroups are non-isomorphic. Note that the right gyrogroups determinned by different k may be isomorphic. For, let k_1 and k_2 be class assigned maps given by $k_1(x) = 0$ for all $x \in S_3$ and $k_2(I) = 0$, $k_2(x) = 6$ for all $x \in S_3 \setminus \{I\}$ respectively. Clearly they are distinct but the corresponding right gyrogroups are the group S_3.

Corollary 3.9. *For $n \geq 3$, there are*

$$\prod_{\sum_{i=1}^{k} n_i \leq n}^{n_i > 1} [n_1, \ldots, n_k]$$

gyrotransversals to the subgroup S_n in the generalized product of group S_n. Here $[n_1, \ldots, n_k]$ denotes the least common multiple of $n_1, \ldots, n_k$.

PROOF: Let $p, q \in S_n$. Since p and p^{-1} both have same cycle structure and p, q are conjugates to each other if and only if they have same cycle structures. Therefore for each $w \in S_n$, the equivalence class $[w]_R$ is same as the conjugacy class of w. Since the class assigned function $k(w)$ has $|w|$ choices on each class $[w]_R$ and the choices are independent on these classes. Therefore the number of class assigned functions k will be $\prod |w|$, where the product runs over each classes with $w \neq 0$. Let $n_1, n_2, \ldots, n_k$ be natural numbers greater than 1 such that $n_1 + n_2 + \ldots + n_k \leq n$. Then order of permutation with cycle structure $(n_1, n_2, \ldots, n_k)$ is $[n_1, n_2, \ldots, n_k]$, the least common multiple of $n_1, n_2, \ldots, n_k$. Thus, by above proposition the number of gyrotransversals to S_n in the generalized product $S_n \times S_n$ will be

$$\prod_{(\sum_{i=1}^{k} n_i) \leq n}^{n_i > 1} [n_1, \ldots, n_k]$$

□

REFERENCES

[1] R. Lal, *Transversals in groups*, Journal of algebra, **181**, 70-81(1996).

[2] R. Lal and A. C. Yadav, *Topological right gyrogroups and gyrotransversals*, Communication in Algebra, 41:9, 3559-3575 (2013).

[3] T. Foguel, A.A. Ungar, *Involutory decomposition of groups into twisted subgroups and subgroups*, J. Group theory, **3**, 27-46 (2000).

[4] T. Foguel, A.A. Ungar, *Gyrogroups and the decomposition of groups into twisted subgroups and subgroups*, Pac. J. of Maths. Vol. 197, No. **1**, (2001).

DEPARTMENT OF MATHEMATICS, M G KASHI VIDYAPITH, VARANASI (INDIA), akhileshyadav538@gmail.com